


Exam # 1 – Physics 203

October 8, 2007

Be sure to include pictures, coordinate systems, etc. where reasonable.

1. For each of the following vectors θ is measured counter clockwise from the positive x-axis. \vec{A} has a magnitude of 4cm and $\theta_A = 30^\circ$, \vec{B} has a magnitude of 7cm and $\theta_B = 112^\circ$.
 - (a) Make a sketch that shows how to graphically add $\vec{A} + \vec{B} = \vec{C}$.
 - (b) Find $\vec{A} + \vec{B} = \vec{C}$ and express \vec{C} using unit vectors.
 - (c) What is the magnitude of \vec{C} .
 - (d) What is the direction of \vec{C} .











2. Two identical billiard balls are labeled A and B . Maryland Fats places ball A at the very edge of the table. He places ball B at the other side. He strikes ball B with his cue so that it flies across the table and off the edge. As it passes A , it just touches ball A lightly, knocking it off. The figure above shows the balls just at the instant they have left the table. Ball B is moving with a speed v_1 and ball A is essentially at rest.

- (a) Which ball do you think will hit the ground first? Explain your reasons for thinking so.

Each graph below shows a quantity vs. time. In each case, the horizontal axis is the time axis. For each of the items below, select which graph could be a plot of that quantity vs. time. If none of the graphs are possible, write N . The time axes are taken to have $t = 0$ at the instant both balls leave the table. Use the x and y axes shown in the figure.

- (b) The x -component of the velocity of ball B .
(c) The y -component of the velocity of ball A .
(d) The y -component of the acceleration of ball A .
(e) The y -component of the force on ball B .
(f) The y -component of the force on ball A .
(g) The x -component of the velocity of ball A .
(h) The y -component of the acceleration of ball B .



3. A slide-loving pig slides down a certain slide (at an angle θ with the ground) in twice the time it would take to slide down a frictionless slide (also at an angle θ with the ground). What is the coefficient of kinetic friction between the pig and the slide? Both slides are the same length.
4. A 5.00kg block is pulled along a horizontal frictionless floor by a cord that exerts a force of magnitude $||\vec{F}|| = 12.0N$ at an angle $\theta = 25^\circ$ above the horizontal. Please ignore frictional effects.
 - (a) What is the magnitude of the block's acceleration?
 - (b) The force magnitude $||\vec{F}||$ is slowly increased. What is its value just before the block is lifted (completely) off the floor?
 - (c) What is the magnitude of the block's acceleration just before it is lifted (completely) off the floor.
5. A contact force between two objects can be thought of as a normal force and a frictional force. Discuss what determines the size and direction of these forces.
6. A wheel of radius 0.30m spins clockwise at 600rpm just above the level ground (the bottom of the wheel just brushes the ground). A chunk of rubber at exactly the 11 o'clock position (this has nothing to do with time, imagine a clock projected on the wheel) breaks free and flies from the wheel.
 - (a) What is the velocity of the chunk just as it breaks free from the wheel?
 - (b) How long is the chunk in the air?
 - (c) What is the velocity of the chunk just before it hits the ground?
 - (d) What is the displacement of the chunk between the time it breaks free and the time it hits the ground?